

1. (7 points) Let  $A, B, C$  be sets. Prove that  $(A - B) \cup (A - C) = A - (B \cap C)$ .

Hint: You may use any one of the following three approaches.

- Write  $(A - B) \cup (A - C) = \{x \in U : p(x)\}$ , where  $p(x) : x \in (A - B) \text{ or } x \in (A - C)$ , and  $A - (B \cap C) = \{x \in U : q(x)\}$ , where  $q(x) : x \in A \text{ and } x \notin (B \cap C)$ , where  $U$  is the universal set. Show that  $p(x)$  and  $q(x)$  are logically equivalent.
- Show that if  $x \in (A - B) \cup (A - C)$ , then  $x \in A - (B \cap C)$ . Also, show that if  $x \in A - (B \cap C)$ , then  $x \in (A - B) \cup (A - C)$ .
- Use set operations such as  $A - X \subseteq A - Y$  if  $Y \subseteq X$ , to argue  $(A - B) \cup (A - C) \subseteq A - (B \cap C)$  and also  $A - (B \cap C) \subseteq (A - B) \cup (A - C)$ .

2. (7 points) Let  $A, B, C$  and  $D$  be sets. Prove that

$$(A \times B) \cap (C \times D) = (A \cap C) \times (B \cap D).$$

Hint: Show that (a) if  $(x, y) \in (A \times B) \cap (C \times D)$ , then  $(x, y) \in (A \cap C) \times (B \cap D)$ , and (b) if  $(x, y) \in (A \cap C) \times (B \cap D)$ , then  $(x, y) \in (A \times B) \cap (C \times D)$ .

3. (7 points) For the following, state whether they are true or not. Then, prove your answer.

- $\forall x \in \mathbf{R}, \exists y \in \mathbf{R}, xy = 1$ ;
- $\exists n \in \mathbf{N}, \exists m \in (\mathbf{N} - \{1\}), nm = 1$ .

Hint: If you want to prove that  $P$  is FALSE, you may try to prove  $\sim P$  is TRUE.

4. (7 points) Show that for any two positive numbers  $a$  and  $b$ ,

$$(a + b) \left( \frac{1}{a} + \frac{1}{b} \right) \geq 4.$$

Hint: Reduce the problem to  $(a + b)^2 \geq 4ab$ , and use algebra.

5. (7 points) Let  $m = 4s + 2$  with  $s \in \mathbf{Z}$ . Show that there are no integers  $x, y$  such that

$$x^2 - y^2 = m.$$

6. (7 points) Prove that the product of an irrational number and a nonzero rational number is irrational.

Hint: Assume that  $x$  is irrational and  $y$  is nonzero rational. If  $xy$  is rational, then ...

7. (8 points) Let  $S = \{a, b, c\} \subseteq \mathbf{Z}$ . For any non-empty subset  $X$  of  $S$ , let  $s(X)$  be the sum of elements in  $X$ . Show that there are non-empty subsets  $A, B$  of  $S$  such that  $s(A) - s(B)$  is divisible by 6.

Hint: For each non-empty subset  $X$  of  $S$ , consider the remainder of  $s(X)$  divided by 6. We get remainders  $r_1, \dots, r_7$ . Show that two of these numbers are the same and deduce the result.

8. (Extra Credit, 8 points) Recall that for a given  $S \subseteq \mathbf{R}$ , the maximum element of  $S$ , denoted by  $\max S$ , is the number  $\alpha \in S$  such that  $\alpha \geq \beta$  for all  $\beta \in S$ .

Let  $A = \{n \in \mathbf{N} : \sqrt{n} \notin \mathbf{Q}\}$ . Show that  $\max A$  does not exist.

Hint: Proof by contradiction. Suppose  $N \in A$  is a maximum. Show that  $n = 2N^2 \in A$  satisfy  $n > N$ .