

SUM OF HERMITIAN MATRICES WITH GIVEN EIGENVALUES: INERTIA, RANK, AND MULTIPLE EIGENVALUES

CHI-KWONG LI* AND YIU-TUNG POON

Abstract

Let A and B be $n \times n$ complex Hermitian (or real symmetric) matrices with eigenvalues $a_1 \geq \cdots \geq a_n$ and $b_1 \geq \cdots \geq b_n$. All possible inertia values, ranks, and multiple eigenvalues of $A + B$ are determined. Extension of the results to the sum of k matrices with $k > 2$, and connections of the results to other subjects such as algebraic combinatorics are also discussed.

2000 Mathematics Subject Classification. 15A42, 15A57.

Key words and phrases. Complex Hermitian matrices, real symmetric matrices, inertia, rank, multiple eigenvalues.

1 Introduction

Let \mathcal{H}_n be the real linear space of $n \times n$ complex Hermitian (or real symmetric) matrices. For a real vector $\mathbf{a} = (a_1, \dots, a_n)$ with $a_1 \geq \cdots \geq a_n$, let

$$\mathcal{H}_n(\mathbf{a}) = \{A \in \mathcal{H}_n : A \text{ has eigenvalues } a_1, \dots, a_n\}.$$

Motivated by problems in pure and applied subjects, there has been a lot of research on the relation between the eigenvalues of $A, B \in \mathcal{H}_n$ and those of $A + B$; [3, 4, 5, 8, 7, 9, 11, 12]. In particular, for given real vectors $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n)$ and $\mathbf{c} = (c_1, \dots, c_n)$ with entries arranged in descending order, a necessary and sufficient condition for the existence of $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ such that $A + B \in \mathcal{H}_n(\mathbf{c})$, or equivalently,

$$\mathcal{H}_n(\mathbf{c}) \subseteq \mathcal{H}_n(\mathbf{a}) + \mathcal{H}_n(\mathbf{b}) \tag{1.1}$$

can be completely described in terms of the equality

$$\sum_{j=1}^n (a_j + b_j - c_j) = 0 \tag{1.2}$$

*Li is an honorary professor of the University of Hong Kong. His research was supported by a USA NSF grant and a HK RCG grant.

and a collection of inequalities in the form

$$\sum_{r \in R} a_r + \sum_{s \in S} b_s \geq \sum_{t \in T} c_t \quad (1.3)$$

for certain m element subsets $R, S, T \subseteq \{1, \dots, n\}$ with $1 \leq m < n$ determined by the Littlewood-Richardson rules; see [5, 7] for details. Using (1.2) and (1.3), we can also deduce the following inequalities

$$\sum_{r \in R^c} a_r + \sum_{s \in S^c} b_s \leq \sum_{t \in T^c} c_t, \quad (1.4)$$

where R^c denotes the complement of R in $\{1, 2, \dots, n\}$. The study has connections to many different areas such as representation theory, algebraic geometry, and algebraic combinatorics, etc.

The set of inequalities in (1.3) grows exponentially with n . Therefore, in spite of the existence of a complete description of the eigenvalues of $A + B$ in terms of those of A and B in \mathcal{H}_n , it is sometimes difficult to answer some basic questions related to the eigenvalues of the matrices A , B and $A + B$. For example, as pointed out by Fulton [7, p.215], given a *proper* subset K of $\{1, 2, \dots, n\}$ and real numbers $\{\gamma_k : k \in K\}$, it is not easy to use the inequalities in (1.3) to determine if there exists \mathbf{c} with $c_k = \gamma_k$ for all $k \in K$ such that (1.1) holds. In particular, the inequalities in (1.3) with $T \subseteq K$ together with those in (1.4) with $T^c \subseteq K$ are necessary but not sufficient for (1.1) in general.

If $K = \{k\}$ is a singleton, then inequalities in (1.3) and (1.4) reduce to the Weyl's inequalities [13] implying that $c_k \in [L_k, R_k]$, where

$$L_k = \max\{a_i + b_j : i + j = n + k\} \quad \text{and} \quad R_k = \min\{a_i + b_j : i + j = k + 1\}. \quad (1.5)$$

Conversely, one can check that for every $c \in [L_k, R_k]$, there exists $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ satisfying $A + B \in \mathcal{H}_n(\mathbf{c})$ with $c_k = c$. So, in this case, the inequalities in (1.3) with $T \subseteq K$ and $c_k = \gamma_k$ for $k \in K$ are also sufficient.

In this paper, we show that if $\mu \in [L_k, L_{k-1}] \cap (R_{k'+1}, R_{k'}]$. Then there exists $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ such that $C = A + B$ has a vector of eigenvalues \mathbf{c} with

$$c_{k-1} < \mu = c_k = c_{k+1} = \dots = c_{k'} < c_{k'+1}.$$

This will follow from a consequence (Corollary 5.7) of the solution of the following problem.

Problem 1.1 *Suppose $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$. Can a given $\mu \in \mathbf{R}$ be an eigenvalue of $A + B$ with a specific multiplicity? Equivalently, can $A + B - \mu I$ have a specific rank?*

We will study the following harder problem.

Problem 1.2 Suppose $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$. Can a given $\mu \in \mathbf{R}$ be an eigenvalue of $A + B$ so that p other eigenvalues are larger than μ , and q other eigenvalues are smaller than μ ? Equivalently, can $A + B - \mu I$ have inertia $(p, q, n - p - q)$, i.e., p positive eigenvalues, q negative eigenvalues, and $n - p - q$ zero eigenvalues?

Clearly, one can replace (A, B) by $(A - \mu I, B)$ and replace $\mathbf{a} = (a_1, \dots, a_n)$ by $(a_1 - \mu, \dots, a_n - \mu)$ so as to focus on the case for $\mu = 0$ in the study.

For two nonnegative integers p and q with $p + q \leq n$, let

$$\mathcal{H}_n(p, q) = \{X \in \mathcal{H}_n : X \text{ has } p \text{ positive eigenvalues and } q \text{ negative eigenvalues}\}.$$

In Section 2, we determine a necessary and sufficient condition on (p, q) for the existence of $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ so that $A + B \in \mathcal{H}_n(p, q)$. In addition, we give a global description of the set of integer pairs (p, q) satisfying these conditions in Section 3. Moreover, we determine those integer pairs (p, q) for the existence of diagonal matrices $A \in \mathcal{H}_n(\mathbf{a})$ and $B \in \mathcal{H}_n(\mathbf{b})$ such that $A + B \in \mathcal{H}_n(p, q)$ in Section 4. Then the results are used to determine all the possible ranks of matrices of the form $A + B$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ in Section 5. We also determine the function $f : \mathbf{R} \rightarrow \mathbf{Z}$ such that $f(\mu)$ is the minimum rank of a matrix of the form $A + B - \mu I$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$. Additional remarks and problems are mentioned in Section 6.

Alternatively, one can describe the results as follows. For $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$, we determine the condition on (p, q) such that $U^*AU + V^*BV \in \mathcal{H}_n(p, q)$ for some unitary matrices U and V , and use the result to determine all possible ranks and multiplicities of eigenvalues of all matrices of the form $U^*AU + V^*BV$.

It turns out that it is more convenient to state and prove the results for $A - B$. We will do this in our discussion and focus on the set

$$\text{In}(\mathbf{a}, \mathbf{b}) = \{(p, q) \in \mathbf{Z} \times \mathbf{Z} : \exists (A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b}) \text{ such that } A - B \in \mathcal{H}_n(p, q)\}.$$

We always assume that $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n)$ and $\mathbf{c} = (c_1, \dots, c_n)$ are real vectors with entries arranged in descending order unless specified otherwise.

2 Characterization of elements in $\text{In}(\mathbf{a}, \mathbf{b})$

First, we obtain an easy to check necessary and sufficient condition for $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$.

Theorem 2.1 Let $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ be real vectors with entries arranged in descending order. Suppose p and q are nonnegative integers satisfying $p + q \leq n$. Then $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ if and only if

(1) $(a_1, \dots, a_{n-q}) - (b_{q+1}, \dots, b_n)$ is a nonnegative vector with at least p positive entries, and

(2) $(b_1, \dots, b_{n-p}) - (a_{p+1}, \dots, a_n)$ is a nonnegative vector with at least q positive entries.

Moreover, if (1) and (2) hold, then there exist block diagonal matrices $A = A_1 \oplus \dots \oplus A_{p+q} \in \mathcal{H}_n(\mathbf{a})$ and $B = B_1 \oplus \dots \oplus B_{p+q} \in \mathcal{H}_n(\mathbf{b})$ with the same block sizes such that $A_j - B_j$ is rank one positive definite for $j = 1, \dots, p$ and $A_j - B_j$ is rank one negative semi-definite for $j = p+1, \dots, p+q$.

Remark 2.2 For fixed $p, q \geq 0$ with $p+q \leq n$, let $K = \{p+1, \dots, n-q\}$. The necessity of condition (1) and (2) in the above theorem can be deduced from the inequalities in (1.3) with $T \subseteq K$ and $c_k = 0$ for $k \in K$. We will give a direct proof of this result for completeness.

It is convenient to use the following notation in our discussion. Suppose $\mathbf{u} = (u_1, \dots, u_m)$ and $\mathbf{v} = (v_1, \dots, v_m)$ are real vectors with entries arranged in descending order. Write $\mathbf{u} \geq_k \mathbf{v}$ (respectively, $\mathbf{u} >_k \mathbf{v}$) if $\mathbf{u} - \mathbf{v}$ is a nonnegative vector with at least (respectively, exactly) k positive entries. We will use $\mathbf{u} \geq \mathbf{v}$ and $\mathbf{u} > \mathbf{v}$ for $\mathbf{u} \geq_0 \mathbf{v}$ and $\mathbf{u} >_n \mathbf{v}$, respectively. For $\mathbf{a} = (a_1, \dots, a_n)$ and $1 \leq m \leq n$, let $\mathbf{a}^m = (a_1, \dots, a_m)$ and $\mathbf{a}_m = (a_{n-m+1}, \dots, a_n)$. One can use these notations to restate conditions (1) and (2) in Theorem 2.1 as

$$\mathbf{a}^{n-q} \geq_p \mathbf{b}_{n-q} \quad \text{and} \quad \mathbf{b}^{n-p} \geq_q \mathbf{a}_{n-p}.$$

The following lemmas are needed to prove Theorem 2.1. The first one was proved in [6].

Lemma 2.3 Let $\tilde{\mathbf{a}} = (\tilde{a}_1, \dots, \tilde{a}_m)$ and $\mathbf{a} = (a_1, \dots, a_n)$ be real vectors with entries arranged in descending order, where $1 \leq m < n$. Then there is $(A, \tilde{A}) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_m(\tilde{\mathbf{a}})$ with \tilde{A} as the leading principal submatrix of A if and only if $a_j \geq \tilde{a}_j \geq a_{n-m+j}$ for $j = 1, \dots, m$.

Lemma 2.4 Let $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$. If $A - B$ is a rank k positive semi-definite matrix, then $\mathbf{a} \geq_k \mathbf{b}$.

Proof. Applying a suitable unitary similarity to $A - B$, we may assume that $A - B = \text{diag}(d_1, \dots, d_k, 0, \dots, 0)$ with $d_1 \geq \dots \geq d_k > 0$. Let $C = B + d_k I_k \oplus 0_{n-k}$ have eigenvalues $c_1 \geq \dots \geq c_n$. Then using the positive semi-definite ordering, we have

$$A \geq C \quad \text{and} \quad B + d_k I \geq C \geq B.$$

By Weyl's inequalities (see [13]), we have

$$a_j \geq c_j \quad \text{and} \quad b_j + d_k \geq c_j \geq b_j, \quad j = 1, \dots, n.$$

Since

$$kd_k = \text{tr}(C - B) = \sum_{j=1}^n (c_j - b_j),$$

and each of the summands on the right side is bounded by d_k , we see that at least k of the summands are positive. It follows that there are at least k indices j such that $a_j > b_j$. \blacksquare

Lemma 2.5 *Let \mathbf{a} and \mathbf{b} be real vectors. Suppose $\{a_1, a_2, \dots, a_n\}$ and $\{b_1, b_2, \dots, b_n\}$ can be partitioned as*

$$\{a_1, a_2, \dots, a_n\} = \bigcup_{j=1}^r \{a_{j,1}, \dots, a_{j,n_j}\} \quad \text{and} \quad \{b_1, b_2, \dots, b_n\} = \bigcup_{j=1}^r \{b_{j,1}, \dots, b_{j,n_j}\}$$

such that for each $1 \leq j \leq r$,

$$a_{j,1} \geq b_{j,1} \geq a_{j,2} \geq b_{j,2} \geq \dots \geq a_{j,n_j} \geq b_{j,n_j}$$

with $a_{j,i} > b_{j,i}$ for at least k_j i 's and $\sum_{j=1}^r k_j \geq m$. Then there exist block diagonal matrices $A = A_1 \oplus \dots \oplus A_m \in \mathcal{H}_n(\mathbf{a})$ and $B = B_1 \oplus \dots \oplus B_m \in \mathcal{H}_n(\mathbf{b})$ with the same block sizes such that $A_j - B_j$ is rank one positive definite for $j = 1, \dots, m$. Consequently, $(m, 0) \in \text{In}(\mathbf{a}, \mathbf{b})$.

Proof. Suppose $r = 1$. We prove the statement by induction on m . When $m = 1$ we have

$$a_1 \geq b_1 \geq a_2 \geq b_2 \geq \dots \geq a_n \geq b_n \tag{2.1}$$

and $a_i > b_i$ for at least one i . If $b_n \geq 0$, then by Lemma 2.4 there is $\tilde{A} \in \mathcal{H}_{n+1}$ with eigenvalues $a_1 \geq \dots \geq a_n \geq a_{n+1} = 0$ such that the leading $n \times n$ submatrix has eigenvalues $b_1 \geq \dots \geq b_n$. Since \tilde{A} is singular, there is $R \in M_n$ and $v \in \mathbf{C}^n$ such that $\tilde{A} = [R|v]^*[R|v]$. Note that $B = RR^*$ and R^*R have the same eigenvalues $b_1 \geq \dots \geq b_n$, and the eigenvalues of $A = [R|v][R|v]^* = RR^* + vv^*$ are the same as the n largest of \tilde{A} and equal to $a_1 \geq \dots \geq a_n$. Thus, there exists unitary $A - B = vv^*$ is rank one positive semi-definite. If $b_n < 0$, apply the argument to $A - b_n I$ and $B - b_n I$ to get the conclusion.

Suppose the result holds for some $m \geq 1$ and (2.1) holds with $a_i > b_i$ for at least $m+1$ i 's. Let $s = \min\{i : a_i > b_i\}$. Then by induction assumption, there exist $A_1, B_1 \in \mathcal{H}_s$ with eigenvalues a_1, \dots, a_s and b_1, \dots, b_s and block diagonal matrices $A_2 \oplus \dots \oplus A_{m+1}$ and $B_2 \oplus \dots \oplus B_{m+1} \in \mathcal{H}_{n-s}$ with eigenvalues a_{s+1}, \dots, a_n and b_{s+1}, \dots, b_n such that $A_j - B_j$ is rank one positive definite for $j = 1, \dots, m+1$. Thus, $A = A_1 \oplus A_2 \oplus \dots \oplus A_{m+1} \in \mathcal{H}_n(\mathbf{a})$ and $B = B_1 \oplus B_2 \oplus \dots \oplus B_{m+1} \in \mathcal{H}_n(\mathbf{b})$ satisfy the requirement.

Now, suppose $r > 1$. Choose non-negative numbers ℓ_j with $\min\{1, k_j\} \leq \ell_j \leq k_j$ for $1 \leq j \leq m$ such that $\ell_1 + \dots + \ell_m = m$. By the result when $r = 1$, there exist block

diagonal matrices A_j and $B_j \in \mathcal{H}_{n_j}$ with eigenvalues $a_{j,1}, \dots, a_{j,n_j}$ and $b_{j,1}, \dots, b_{j,n_j}$ such that $A_j - B_j$ is positive semi-definite with rank ℓ_j . Thus, for $A = A_1 \oplus \dots \oplus A_m$ and $B = B_1 \oplus \dots \oplus B_m$, $A - B$ is positive semi-definite with rank m . \blacksquare

We are now ready to present the following.

Proof of Theorem 2.1. Suppose $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ satisfies $A - B \in \mathcal{H}_n(p, q)$. Applying a unitary similarity to $A - B$, we may assume that $A - B = \text{diag}(c_1, \dots, c_n)$ such that $c_1 \geq \dots \geq c_p > 0 = c_{p+1} = \dots = c_{n-q} = 0 > c_{n-q+1} \geq \dots \geq c_n$. Let

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$$

with $A_{11}, B_{11} \in \mathcal{H}_{n-q}$. Then $A_{11} - B_{11}$ is positive semi-definite with p positive eigenvalues. Suppose A_{11} and B_{11} have eigenvalues $\alpha_1 \geq \dots \geq \alpha_{n-q}$ and $\beta_1 \geq \dots \geq \beta_{n-q}$, respectively. By Lemmas 2.3 and 2.4, we have

$$(a_1, \dots, a_{n-q}) \geq (\alpha_1, \dots, \alpha_{n-q}) \geq_p (\beta_1, \dots, \beta_{n-q}) \geq (b_{q+1}, \dots, b_n)$$

This proves (1). Similarly, we can prove condition (2).

To prove the converse, given real vectors \mathbf{a} and \mathbf{b} , we first show that for every n , the result holds if $pq = 0$ or $p + q = n$. If $(p, q) = (0, 0)$, then we have $\mathbf{a} = \mathbf{b}$ and the result follows.

Suppose $p > 0$ and $q = 0$. Let $n = rp + s$, with $r \geq 0$ and $1 \leq s \leq p$ (not $0 \leq s < p$ as given by the Euclidean algorithm). Then (1) and (2) imply that

$$a_i \geq b_i \geq a_{p+i} \geq \dots \geq a_{rp+i} \geq b_{rp+i} \quad \text{for } 1 \leq i \leq s$$

$$a_j \geq b_j \geq a_{p+j} \geq \dots \geq a_{(r-1)p+j} \geq b_{(r-1)p+j} \quad \text{for } s+1 \leq j \leq p$$

with $a_i > b_i$ for at least p i 's. Therefore the result follows from Lemma 2.5.

Similarly, the result holds for $p = 0$ and $q > 0$. Hence, the result holds if $pq = 0$.

For $p + q = n$, Let $A = \text{diag}(a_1, \dots, a_n)$ and $B = \text{diag}(b_{q+1}, \dots, b_n, b_1, \dots, b_{n-p})$. Then it follows from (1) and (2) that $A - B \in \mathcal{H}_n(p, q)$.

We complete the proof of the converse by induction on n . The result is clear for $n \leq 2$.

Assume that the result is valid for all real vectors of lengths less than n . Suppose $(p, q) \geq (1, 1)$, $p + q < n$, and the inequalities in (1) and (2) hold. Then we have

$$a_i \geq b_{q+i} \quad \text{for } 1 \leq i \leq n - q \tag{2.2}$$

and

$$b_i \geq a_{p+i} \quad \text{for } 1 \leq i \leq n - p \tag{2.3}$$

with at least p strict inequalities hold in (2.2) and at least q strict inequalities hold in (2.3).

If $a_i = b_{q+i}$ for some $1 \leq i \leq n - q$, then letting $\mathbf{a}' = (a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n)$ and $\mathbf{b}' = (b_1, \dots, b_{q+i-1}, b_{q+i+1}, \dots, b_n)$, we have

$$\begin{aligned} 1 \leq j < i &\Rightarrow a'_j = a_j \geq b_{q+j} = b'_{q+j} \\ i \leq j \leq n - 1 - q &\Rightarrow a'_j = a_{j+1} \geq b_{q+j+1} = b'_{q+j} \end{aligned} \quad (2.4)$$

$$\begin{aligned} 1 \leq j < i - p &\Rightarrow b'_j = b_j \geq a_{p+j} = a'_{p+j} \\ i - p \leq j < i + q &\Rightarrow b'_j = b_j \geq a_{p+j} \geq a_{p+j+1} = a'_{p+j} \\ i + q \leq j \leq n - 1 - p &\Rightarrow b'_j = b_{j+1} \geq a_{p+j+1} = a'_{p+j} \end{aligned} \quad (2.5)$$

with at least p strict inequalities hold in (2.4) and at least q strict inequalities hold in (2.5). By induction hypothesis, there exist $A', B' \in \mathcal{H}_{n-1}$ with eigenvalues $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n$ and $b_1, \dots, b_{q+i-1}, b_{q+i+1}, \dots, b_n$ such that $A' - B' \in \mathcal{H}_{n-1}(p, q)$. Hence, $[a_i] \oplus A' - [b_{q+i}] \oplus B' \in \mathcal{H}_n(p, q)$.

Similarly, the result holds if $b_i = a_{p+i}$ for some $1 \leq i \leq n - p$.

So, we may assume that all inequalities are strict in (2.2) and (2.3). By symmetry, we may assume that $q \leq p$. Since $n > p + q$, let $n = r(p + q) + s$, where $r > 0$ and $1 \leq s \leq p + q$. We will arrange a_1, \dots, a_n and b_1, \dots, b_n in $p + q$ chains of inequalities so that Lemma 2.5 can be applied. To this end, define $m = \min\{s, q, p + q - s\}$,

$$i_1 = \max\{1, s - q + 1\}, \quad i_2 = \min\{s, p\}, \quad j_1 = \max\{1, s - p + 1\}, \quad \text{and} \quad j_2 = \min\{s, q\}.$$

We have

	$1 \leq s \leq q$	$q < s \leq p$	$p < s \leq p + q$
$i_1 = \max\{1, s - q + 1\}$	1	$s - q + 1$	$s - q + 1$
$i_2 = \min\{s, p\}$	s	s	p
$j_1 = \max\{1, s - p + 1\}$	1	1	$s - p + 1$
$j_2 = \min\{s, q\}$	s	q	q
$m = \min\{s, q, p + q - s\}$	s	q	$p + q - s$

Then $i_2 - i_1 = j_2 - j_1 = m - 1$. By conditions (1) and (2), we can list all the entries of \mathbf{a} and \mathbf{b} in the following $p + q$ chains of interlacing inequalities:

$$\begin{array}{cccccccc} a_1 & > & b_{q+1} & > & a_{p+q+1} & > & \cdots & > & b_{(r-1)(p+q)+q+1} & > & a_{r(p+q)+1} & > & b_{r(p+q)+q+1} \\ \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots \\ a_{i_1-1} & > & b_{q+i_1-1} & > & a_{p+q+i_1-1} & > & \cdots & > & b_{(r-1)(p+q)+q+i_1-1} & > & a_{r(p+q)+i_1-1} & > & b_{r(p+q)+q+i_1-1} \\ a_{i_1} & > & b_{q+i_1} & > & a_{p+q+i_1} & > & \cdots & > & b_{(r-1)(p+q)+q+i_1} & > & a_{r(p+q)+i_1} & > & \\ \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots \\ a_{i_2} & > & b_{q+i_2} & > & a_{p+q+i_2} & > & \cdots & > & b_{(r-1)(p+q)+q+i_2} & > & a_{r(p+q)+i_2} & > & \\ a_{i_2+1} & > & b_{q+i_2+1} & > & a_{p+q+i_2+1} & > & \cdots & > & b_{(r-1)(p+q)+q+i_2+1} & > & & > & \\ \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots \\ a_p & > & b_{q+p} & > & a_{p+q+p} & > & \cdots & > & b_{r(p+q)}, & & & & \end{array}$$

and

$$\begin{array}{cccccccc}
b_1 & > & a_{p+1} & > & b_{p+q+1} & > & \cdots & > & a_{(r-1)(p+q)+p+1} & > & b_{r(p+q)+1} & > & a_{r(p+q)+p+1} \\
\vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots \\
b_{j_1-1} & > & a_{p+j_1-1} & > & b_{p+q+j_1-1} & > & \cdots & > & a_{(r-1)(p+q)+p+j_1-1} & > & b_{r(p+q)+j_1-1} & > & a_{r(p+q)+p+j_1-1} \\
b_{j_1} & > & a_{p+j_1} & > & b_{p+q+j_1} & > & \cdots & > & a_{(r-1)(p+q)+p+j_1} & > & b_{r(p+q)+j_1} & > & a_{r(p+q)+p+j_1} \\
\vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots \\
b_{j_2} & > & a_{p+j_2} & > & b_{p+q+j_2} & > & \cdots & > & a_{(r-1)(p+q)+p+j_2} & > & b_{r(p+q)+j_2} & > & a_{r(p+q)+p+j_2} \\
b_{j_2+1} & > & a_{p+j_2+1} & > & b_{p+q+j_2+1} & > & \cdots & > & a_{(r-1)(p+q)+p+j_2+1} & > & b_{r(p+q)+j_2+1} & > & a_{r(p+q)+p+j_2+1} \\
\vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots & > & \vdots \\
b_q & > & a_{p+q} & > & b_{p+q+q} & > & \cdots & > & a_{r(p+q)} & > & b_{r(p+q)} & > & a_{r(p+q)}
\end{array}$$

where a_i and b_i would not appear if $i < 0$ or $i > n$.

In fact, it is easy to construct the p chains of inequalities in the first list and q chains of inequalities in the second list as follows. Put the first p entries of \mathbf{a} vertically in the first column of the first list, the next q entries of \mathbf{a} vertically in the second column of the second list, then the next p entries of \mathbf{a} in the third column of first list, and so forth. Similarly, put the first q entries of \mathbf{b} in the first column of the second list, the next p entries of \mathbf{b} in the second column of the first list, then the next q entries of \mathbf{b} in the third column of the second list, and so forth.

For the application of Lemma 2.5, the chains of inequalities with starting terms a_i for $i_1 \leq i \leq i_2$ are not acceptable because the first and last terms are entries of \mathbf{a} . Similarly, the chains of inequalities with starting terms b_j for $j_1 \leq j \leq j_2$ are not acceptable. Since $i_2 - i_1 = j_2 - j_1$, we can amend the situations as follows. For $i_1 \leq i \leq i_2$, let $i' = j_1 + i - i_1$. Then $j_1 \leq i' \leq j_2$ and we can replace the pair of interlacing inequalities

$$\begin{array}{cccccccc}
a_i & > & b_{q+i} & > & a_{p+q+i} & > & \cdots & > & b_{(r-1)(p+q)+q+i} & > & a_{r(p+q)+i}, \\
b_{i'} & > & a_{p+i'} & > & b_{p+q+i'} & > & \cdots & > & a_{(r-1)(p+q)+p+i'} & > & b_{r(p+q)+i'},
\end{array}$$

by one of the following pairs:

$$\begin{array}{cccccccc}
a_i & > & b_{q+i} & > & a_{p+q+i} & > & \cdots & > & b_{(r-1)(p+q)+q+i} & > & a_{r(p+q)+i} & > & b_{r(p+q)+i'}, \\
b_{i'} & > & a_{p+i'} & > & b_{p+q+i'} & > & \cdots & > & a_{(r-1)(p+q)+p+i'} & > & a_{r(p+q)+i} & > & b_{r(p+q)+i'},
\end{array}$$

if $a_{r(p+q)+i} > b_{r(p+q)+i'}$, or

$$\begin{array}{cccccccc}
a_i & > & b_{q+i} & > & a_{p+q+i} & > & \cdots & > & b_{(r-1)(p+q)+q+i}, \\
b_{i'} & > & a_{p+i'} & > & b_{p+q+i'} & > & \cdots & > & a_{(r-1)(p+q)+p+i'} & > & b_{r(p+q)+i'} & \geq & a_{r(p+q)+i},
\end{array}$$

if $a_{r(p+q)+i} \leq b_{r(p+q)+i'}$. After the above modification, we can apply Lemma 2.5 to the eigenvalues in the interlacing inequalities with starting terms a_i to get a rank p positive semi-definite matrix, and then apply Lemma 2.5 to the eigenvalues in the interlacing inequalities with starting terms b_j to get a rank q semi-definite matrix. The result follows.

Following our proof, one can construct the matrices A and B in block diagonal forms as asserted in the last statement of the theorem. \blacksquare

It is easy to use Theorem 2.1 to test whether a given pair of integers (p, q) belongs to $\text{In}(\mathbf{a}, \mathbf{b})$. Here is an example.

Example 2.6 Let $\mathbf{a} = (6, 6, 4, 3, 3, 2, 1)$ and $\mathbf{b} = (5, 4, 3, 3, 1, 1, 1)$. Then the following hold.

(a) $(1, 1) \notin \text{In}(\mathbf{a}, \mathbf{b})$ as $(b_1, \dots, b_{7-1}) - (a_{1+1}, \dots, a_7) = (5, 4, 3, 3, 1, 1) - (6, 4, 3, 3, 2, 1)$ has a negative entry.

(b) $(2, 0) \in \text{In}(\mathbf{a}, \mathbf{b})$ as $(a_1, \dots, a_{7-0}) - (b_{1+0}, \dots, b_7) = (6, 6, 4, 3, 3, 2, 1) - (5, 4, 3, 3, 1, 1, 1) = (1, 2, 1, 0, 2, 1, 0)$ and $(b_1, \dots, b_{7-2}) - (a_{2+1}, \dots, a_7) = (5, 4, 3, 3, 1) - (4, 3, 3, 2, 1) = (1, 1, 0, 1, 0)$.


In fact, if $A = \text{diag}(6, 4, 6, 2, 3, 3, 1)$ and $B = B_1 \oplus B_2$ with

$$B_1 = \begin{pmatrix} 7/2 & \sqrt{15}/2 \\ \sqrt{15}/2 & 5/2 \end{pmatrix} \quad \text{and} \quad B_2 = \begin{pmatrix} 7/2 & \sqrt{5}/2 \\ \sqrt{5}/2 & 3/2 \end{pmatrix} \oplus \text{diag}(3, 3, 1),$$

then $(A, B) \in \mathcal{H}_7(\mathbf{a}) \times \mathcal{H}_7(\mathbf{b})$ such that

$$A - B = \begin{pmatrix} 5/2 & -\sqrt{15}/2 \\ -\sqrt{15}/2 & 3/2 \end{pmatrix} \oplus \left[\begin{pmatrix} 5/2 & -\sqrt{5}/2 \\ -\sqrt{5}/2 & 1/2 \end{pmatrix} \oplus \text{diag}(0, 0, 0) \right] \in \mathcal{H}_7(2, 0).$$

We can also test every (p, q) pair of nonnegative integers with $p + q \leq 7$ and depict the set $\text{In}(\mathbf{a}, \mathbf{b})$ as points in \mathbf{R}^2 as follows.



Corollary 2.7 Suppose $(p_1, q_1), (p_2, q_2) \in \text{In}(\mathbf{a}, \mathbf{b})$. Let $p = \min\{p_1, p_2\}$ and $q = \min\{q_1, q_2\}$. Then $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$.

Proof. Suppose $p = p_i$ and $q = q_j$. Since $(p_1, q_1), (p_2, q_2) \in \text{In}(\mathbf{a}, \mathbf{b})$, we have

$$\begin{aligned} \mathbf{a}^{n-q_j} \geq_{p_j} \mathbf{b}_{n-q_j} &\Rightarrow \mathbf{a}^{n-q} \geq_p \mathbf{b}_{n-q} && \text{and} \\ \mathbf{b}^{n-p_i} \geq_{q_i} \mathbf{a}_{n-p_i} &\Rightarrow \mathbf{b}^{n-p} \geq_q \mathbf{a}_{n-p}. \end{aligned}$$

Hence, by Theorem 2.1, $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$. ■

3 A global description of $\text{In}(\mathbf{a}, \mathbf{b})$

While Theorem 2.1 allows us to test if a pair of nonnegative integers lies in $\text{In}(\mathbf{a}, \mathbf{b})$, it would be nice to have a global description of the region for all integer pairs in $\text{In}(\mathbf{a}, \mathbf{b})$. The objective of this section is to obtain such a description.

Note that if \mathbf{a} and \mathbf{b} has a common entry with multiplicities n_1 and n_2 in the two vectors such that $n_1 + n_2 > n$, then for any $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$, the null space of $A - B$ has dimension at least $n_1 + n_2 - m$, and a reduction of the vectors \mathbf{a} and \mathbf{b} is possible in the problem of describing $\text{In}(\mathbf{a}, \mathbf{b})$ as shown in the following proposition.

Proposition 3.1 *Let $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$, be two real vectors with entries arranged in descending order. Suppose $a_i = a_{i+1} = \dots = a_{i+n_1-1} = b_j = b_{j+1} = \dots = b_{j+n_2-1}$, for some $i, j, n_1, n_2 \geq 1$ such that $n_1 + n_2 > n$. Let $s = n_1 + n_2 - n$ and \mathbf{a}' , \mathbf{b}' be obtained by deleting s a_i from each of \mathbf{a} and \mathbf{b} . Then $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ if and only if $(p, q) \in \text{In}(\mathbf{a}', \mathbf{b}')$.*

Proof. Suppose A and B have eigenvalues a_1, \dots, a_n and b_1, \dots, b_n . Then the intersection of the eigenspaces of A and B associated with a_i has dimension $\geq s$. So there exists a unitary U such that $U^*AU = A' \oplus a_i I_s$ and $U^*BU = B' \oplus a_i I_s$. Therefore, $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ if and only if $(p, q) \in \text{In}(\mathbf{a}', \mathbf{b}')$. ■

By the above lemma, to describe $\text{In}(\mathbf{a}, \mathbf{b})$, we can focus on the (\mathbf{a}, \mathbf{b}) pair such that \mathbf{a} and \mathbf{b} do not have a common entry whose multiplicities in the two vectors have sum exceeding n . To describe the main result in this section, we need the following definition.

Definition 3.2 *Suppose $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ are real vectors with entries arranged in descending order. Let*

$$p_0 = \begin{cases} n & \text{if } b_1 < a_n, \\ \min\{t : 0 \leq t < n, \mathbf{b}^{n-t} \geq \mathbf{a}_{n-t}\} & \text{otherwise;} \end{cases} \quad (3.1)$$

$$q_0 = \begin{cases} n & \text{if } a_1 < b_n, \\ \min\{t : 0 \leq t < n, \mathbf{a}^{n-t} \geq \mathbf{b}_{n-t}\} & \text{otherwise.} \end{cases} \quad (3.2)$$

Suppose

$$(a_1, \dots, a_n, b_1, \dots, b_n) \text{ has no entry with multiplicity larger than } n. \quad (3.3)$$

Let

$$k = \begin{cases} n - p_0 & \text{if } b_1 \leq a_n, \\ \min\{t : 0 \leq t < n - p_0, \mathbf{b}^{n-p_0-t} > \mathbf{a}_{n-p_0-t}\} & \text{otherwise;} \end{cases} \quad (3.4)$$

$$\ell = \begin{cases} n - q_0 & \text{if } a_1 \leq b_n, \\ \min\{t : 0 \leq t < n - q_0, \mathbf{a}^{n-q_0-t} > \mathbf{b}_{n-q_0-t}\} & \text{otherwise.} \end{cases} \quad (3.5)$$

Furthermore, for $0 \leq i \leq n - (p_0 + q_0 + \ell)$ and $0 \leq j \leq n - (p_0 + q_0 + k)$, let

$$Q_i \text{ be the number of positive entries in } \mathbf{b}^{n-p_i} - \mathbf{a}_{n-p_i} \quad \text{with } p_i = p_0 + i, \quad (3.6)$$

$$P_j \text{ be the number of positive entries in } \mathbf{a}^{n-q_j} - \mathbf{b}_{n-q_j} \quad \text{with } q_j = q_0 + j. \quad (3.7)$$

In Example 2.6, we have $(k, \ell) = (1, 1)$,

$$\begin{aligned} (p_0, q_0) &= (2, 0), (p_0, Q_0) = (2, 3), (P_0, q_0) = (5, 0), \\ (p_1, Q_1) &= (3, 4) = (P_4, q_4), (p_2, Q_2) = (4, 3) = (P_3, q_3), \\ (p_3, Q_3) &= (5, 2) = (P_2, q_2), (p_4, Q_4) = (6, 1) = (P_1, q_1). \end{aligned}$$

In general, we will show in Lemma 3.11 that $p_k \leq P_\ell$ and $p_i + Q_i = n = P_j + q_j$ for all $k \leq i \leq n - (p_0 + q_0 + \ell)$ and $\ell \leq j \leq n - (p_0 + q_0 + k)$. Therefore, the points in

$$\{(p_i, Q_i) : k \leq i \leq n - (p_0 + q_0 + \ell)\} \cup \{(P_j, q_j) : \ell \leq j \leq n - (p_0 + q_0 + k)\}$$

lie on the line segment joining (p_k, Q_k) and (P_ℓ, q_ℓ) .

Theorem 3.3 *Let \mathbf{a} and \mathbf{b} be real vectors satisfying condition (3.3). Use the notation in Definition 3.2. The following conditions hold.*

(1) *The polygon \mathcal{P} obtained by joining the points*

$$(p_0, q_0), (p_0, Q_0), (p_1, Q_1), \dots, (p_k, Q_k), (P_\ell, q_\ell), (P_{\ell-1}, q_{\ell-1}), \dots, (P_0, q_0), (p_0, q_0)$$

is convex.

(2) *$\text{In}(\mathbf{a}, \mathbf{b})$ consists of all the integer pairs (p, q) in \mathcal{P} .*

In Example 2.6, \mathcal{P} is obtained by joining $(2, 0), (2, 3), (3, 4), (6, 1), (5, 0), (2, 0)$. Before presenting the proof of the theorem, we illustrate how to use the theorem in the following corollaries.

Corollary 3.4 *Suppose \mathbf{a} and \mathbf{b} be real vectors with no common entries. Using the notation in (3.1) and (3.2), we have*

$$\text{In}(\mathbf{a}, \mathbf{b}) = \{(p, q) : p \geq p_0, q \geq q_0, p + q \leq n\}.$$

Proof. Since \mathbf{a} and \mathbf{b} have no common entries, we see that for each $i \in \{1, \dots, k\}$, the vector $\mathbf{b}^{n-p_i} - \mathbf{a}_{n-p_i}$ is positive, and hence $p_i + Q_i = n$. Similarly, $P_j + q_j = n$ for each $j \in \{1, \dots, \ell\}$. By Theorem 3.3, the result follows. ■

Corollary 3.5 *Suppose there are $\mu > \nu$ and $0 \leq u, v \leq n$ such that*

$$\mu = a_1 = \cdots = a_u = b_1 = \cdots = b_\nu \quad \text{and} \quad \nu = a_{u+1} = \cdots = a_n = b_{\nu+1} = \cdots = b_n,$$

then

$$\text{In}(\mathbf{a}, \mathbf{b}) = \{(u - w, v - w) : \max\{0, u + v - n\} \leq w \leq \min\{u, v\}\}.$$

Proof. Without loss of generality, we may assume that $u \geq v$, $\mu = 1$ and $\nu = 0$. Furthermore, by Proposition 3.1, we may assume that $u + v = n$. Then $(p_0, q_0) = (u - v, 0)$. Moreover, $(p_i, Q_i) = (p_0 + i, i) = (P_i, q_i)$ for $i = 1, \dots, v$. By Theorem 3.3, the result follows. \blacksquare

We establish some lemmas to prove Theorem 3.3. The first three lemmas give additional properties of p_0, q_0, P_i, Q_j , and confirm that $(p_0, q_0), (p_i, Q_i), (P_j, q_j) \in \text{In}(\mathbf{a}, \mathbf{b})$.

Lemma 3.6 *Suppose \mathbf{a}, \mathbf{b} are two real vectors, and p_0, q_0 are defined by (3.1) and (3.2). Then the following conditions hold.*

- (1) $p_0 = \min\{p : (p, q) \in \text{In}(\mathbf{a}, \mathbf{b}) \text{ for some } q \geq 0\}$, and $\mathbf{a}^{p_0} - \mathbf{b}_{p_0}$ is a positive vector if $p_0 > 0$.
- (2) $q_0 = \min\{q : (p, q) \in \text{In}(\mathbf{a}, \mathbf{b}) \text{ for some } p \geq 0\}$, and $\mathbf{b}^{q_0} - \mathbf{a}_{q_0}$ is a positive vector if $q_0 > 0$.
- (3) $(p_0, q_0) \in \text{In}(\mathbf{a}, \mathbf{b})$.

Proof. (1) Suppose p_0 is given by (3.1). If $p_0 = n$, then $b_1 < a_n$ and $\text{In}(\mathbf{a}, \mathbf{b}) = \{(n, 0)\}$. If $p_0 < n$, then we have $b_j \geq a_{p_0+j}$ for all $1 \leq j \leq n - p_0$. Let $A = \text{diag}(a_1, \dots, a_n)$ and $B = \text{diag}(b_{n-p_0+1}, \dots, b_n, b_1, \dots, b_{n-p_0})$. Then $A - B$ has at most p_0 positive eigenvalues. Therefore,

$$p_0 \geq \min\{p : (p, q) \in \text{In}(\mathbf{a}, \mathbf{b}) \text{ for some } q \geq 0\}.$$

On the other hand, suppose $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ for some $q \geq 0$. Then there exists $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ such that $A - B \in \mathcal{H}_n(p, q)$. By Theorem 2.1, we have $\mathbf{b}^{n-p} \geq \mathbf{a}_{n-p}$. Therefore, $p \geq p_0$. Hence,

$$p_0 \leq \min\{p : (p, q) \in \text{In}(\mathbf{a}, \mathbf{b}) \text{ for some } q \geq 0\}.$$

If $p_0 > 0$, then there exists $1 \leq i \leq n - (p_0 - 1)$ such that $a_{p_0-1+i} > b_i$. So, for all $1 \leq j \leq p_0$, we have

$$a_j \geq a_{p_0-1+i} > b_i \geq b_{n-p_0+j}$$

i.e., $\mathbf{a}^{p_0} - \mathbf{b}_{p_0}$ is positive. This proves (1). The proof of (2) is similar.

(3) By the results in (1) and (2), we can choose $p \geq p_0$ and $q \geq q_0$ such that (p, q) and $(p_0, q) \in \text{In}(\mathbf{a}, \mathbf{b})$. Hence, by Corollary 2.7, $(p_0, q_0) \in \text{In}(\mathbf{a}, \mathbf{b})$. \blacksquare

Note that assumption (3.3) is not needed in Lemma 3.6.

Lemma 3.7 Suppose \mathbf{a} and \mathbf{b} are real vectors satisfying condition (3.3). Let $s \in \{0, \dots, n-1\}$ be such that $\mathbf{b}^{n-s} - \mathbf{a}_{n-s}$ has a non-positive entry. Then $\mathbf{a}^{s+1} - \mathbf{b}_{s+1}$ is positive.

Proof. Suppose the conclusion is not true. Then $\mathbf{a}^{s+1} - \mathbf{b}_{s+1}$ is not positive. Hence there is $i \in \{1, \dots, s+1\}$ such that $a_i \leq b_{n-s-1+i}$. Since the vector $\mathbf{b}^{n-s} - \mathbf{a}_{n-s}$ has a non-positive entry, $b_j \leq a_{s+j}$ for some $j \in \{1, \dots, n-s\}$. Hence

$$b_j \leq a_{s+j} \leq a_{s+j-1} \leq \dots \leq a_i \leq b_{n-s-1+i} \leq \dots \leq b_j.$$

Consequently, all the inequalities become equalities, and the multiplicity of $a_i = b_j$ in the vector $(a_1, \dots, a_n, b_1, \dots, b_n)$ equals $(s+j-i+1) + (n-s+i-j) = n+1$, contradicting assumption (3.3). \blacksquare

By Lemma 3.7 and the definition of ℓ and k , we see that $(n-q_0-\ell, q_0+\ell), (n-p_0-k, p_0+k) \in \text{In}(\mathbf{a}, \mathbf{b})$ if \mathbf{a}, \mathbf{b} satisfy (3.3).

Lemma 3.8 Let \mathbf{a} and \mathbf{b} be real vectors satisfying (3.3). Use the notation in Definition 3.2. For $0 \leq i \leq n - (p_0 + q_0 + \ell)$ and $0 \leq j \leq n - (p_0 + q_0 + k)$, we have

- (1) $\mathbf{a}^{p_0+i} > \mathbf{b}_{p_0+i}$ and $\mathbf{b}^{q_0+j} > \mathbf{a}_{q_0+j}$.
- (2) $(p_i, Q_i), (P_j, q_j) \in \text{In}(\mathbf{a}, \mathbf{b})$.
- (3) $Q_i = \max\{q : (p_0 + i, q) \in \text{In}(\mathbf{a}, \mathbf{b})\}$ and $P_j = \max\{p : (p, q_0 + j) \in \text{In}(\mathbf{a}, \mathbf{b})\}$.
- (4) $p_0 + q_0 + k + \ell \leq n$.

Proof. If p_0 or $q_0 = n$, then $k = \ell = 0$, and the results follow. Therefore, in the rest of the proof, we assume that $p_0, q_0 < n$.

(1) It follows from the definition of ℓ and k that $\mathbf{a}^{n-q_0-\ell} > \mathbf{b}_{n-q_0-\ell}$ and $\mathbf{b}^{n-p_0-k} > \mathbf{a}_{n-p_0-k}$. For $0 \leq i \leq n - (p_0 + q_0 + \ell)$, we have $p_0 + i \leq n - q_0 - \ell$. Therefore, $\mathbf{a}^{p_0+i} > \mathbf{b}_{p_0+i}$. Similarly, $\mathbf{b}^{q_0+j} > \mathbf{a}_{q_0+j}$ for $0 \leq j \leq n - (p_0 + q_0 + k)$.

(2) Since, $\text{diag}(a_1, \dots, a_n) - \text{diag}(b_{n-p_i+1}, \dots, b_n, b_1, \dots, b_{n-p_i}) \in \mathcal{H}_n(p_i, Q_i)$, we have $(p_i, Q_i) \in \text{In}(\mathbf{a}, \mathbf{b})$. Similarly, $(P_j, q_j) \in \text{In}(\mathbf{a}, \mathbf{b})$.

(3) Suppose $(p_i, q) \in \text{In}(\mathbf{a}, \mathbf{b})$. Then $\mathbf{b}^{n-p_i} \geq_q \mathbf{a}_{n-p_i}$. So, $q \leq Q_i$. Hence,

$$Q_i = \max\{q : (p_0 + i, q) \in \text{In}(\mathbf{a}, \mathbf{b})\}.$$

Similarly, we have

$$P_j = \max\{p : (p, q_0 + j) \in \text{In}(\mathbf{a}, \mathbf{b})\}.$$

(4) Since $(n-q_0-\ell, q_0+\ell) \in \text{In}(\mathbf{a}, \mathbf{b})$, we have $n-q_0-\ell \geq p_0$ by Lemma 3.6. From the definition of k and $\mathbf{a}^{n-q_0-\ell} > \mathbf{b}_{n-q_0-\ell}$, we have $n-q_0-\ell \geq p_0+k$. Thus, $p_0+q_0+k+\ell \leq n$. \blacksquare

Clearly, P_j is equal to $n - q_j$ minus the number of zero entries in $\mathbf{a}^{n-q_j} - \mathbf{b}_{n-q_j}$. Therefore, in order to study the relationship between P_j and P_{j+1} , we need to keep track of the zero entries in the vector $\mathbf{a}^{n-q_j} - \mathbf{b}_{n-q_j}$ and investigate how they are related to the entries of $\mathbf{a}^{n-q_{j-1}} - \mathbf{b}_{n-q_{j-1}}$. For this reason, we introduce the following definition.

Definition 3.9 For $1 \leq i \leq j \leq m \leq n$, we say that $[i, j] = \{t : i \leq t \leq j\}$ is a maximal interval of $(\mathbf{a}^m, \mathbf{b}_m)$ if

$$\begin{aligned} a_{i-1} &> a_i &= a_{i+1} &= \cdots = a_j \\ &= b_{n-m+i} &= b_{n-m+i+1} &= \cdots = b_{n-m+j} > b_{n-m+j+1}. \end{aligned}$$

The length of a maximal interval $[i, j]$ is given by $j - i + 1$. The set of all maximal interval of $(\mathbf{a}^m, \mathbf{b}_m)$ will be denoted by $S(\mathbf{a}^m, \mathbf{b}_m)$. Let $T = T(\mathbf{a}^m, \mathbf{b}_m)$ be the maximum length of a maximal interval of $(\mathbf{a}^m, \mathbf{b}_m)$. For $1 \leq t \leq T$, let s_t be the number of maximal intervals of $(\mathbf{a}^m, \mathbf{b}_m)$ with length t . The sequence (s_1, s_2, \dots, s_T) will be denoted by $\mathbf{s}(\mathbf{a}^m, \mathbf{b}_m)$.

Lemma 3.10 Suppose $\mathbf{a}^m \geq \mathbf{b}_m$ for some $1 \leq m \leq n$. Then the following conditions hold.

- (1) $\mathbf{a}^m >_q \mathbf{b}_m$ where $q = m - \sum_{t=1}^T t s_t$.
- (2) $[i, j] \in S(\mathbf{a}^{m-1}, \mathbf{b}_{m-1})$ if and only if $[i, j+1] \in S(\mathbf{a}^m, \mathbf{b}_m)$.
- (3) $\mathbf{a}^{m-1} >_{q_1} \mathbf{b}_{m-1}$, where $q_1 = q - 1 + \sum_{t=1}^T s_t$.
- (4) If $\mathbf{a}^{m-2} >_{q_2} \mathbf{b}_{m-2}$, then $q_2 - q_1 \leq q_1 - q$.

Here, we assume that $m > 1$ for (2) – (3) and $m > 2$ for (4).

Proof. Condition (1) holds because $\sum_{t=1}^T t s_t$ is the number of zero entries in $\mathbf{a}^m - \mathbf{b}_m$. To prove (2), suppose $[i, j] \in S(\mathbf{a}^{m-1}, \mathbf{b}_{m-1})$. Then we have

$$\begin{aligned} a_{i-1} &> a_i &= a_{i+1} &= \cdots = a_j \\ &= b_{n-(m-1)+i} &= b_{n-(m-1)+i+1} &= \cdots = b_{n-(m-1)+j} > b_{n-(m-1)+j+1}. \end{aligned} \tag{3.8}$$

Since $a_{i-1} > a_i \geq b_{n-m+i} \geq b_{n-m+i+1} = a_i$ and $a_j \geq a_{j+1} \geq b_{n-m+j+1} = a_j > b_{n-(m-1)+j+1}$, we have $a_i = b_{n-m+i} = b_{n-m+i+1}$ and $a_j = a_{j+1} = b_{n-m+j+1}$. This gives

$$\begin{aligned} a_{i-1} &> a_i &= a_{i+1} &= \cdots = a_{j+1} \\ &= b_{n-m+i} &= b_{n-m+i+1} &= \cdots = b_{n-m+j+1} > b_{n-m+j+2}. \end{aligned} \tag{3.9}$$

Thus, $[i, j+1] \in S(\mathbf{a}^m, \mathbf{b}_m)$. Conversely, if $[i, j+1] \in S(\mathbf{a}^m, \mathbf{b}_m)$ for some $j \geq i$, then (3.9) holds. Thus (3.8) follows and $[i, j] \in S(\mathbf{a}^{m-1}, \mathbf{b}_{m-1})$.

To prove (3), let $\mathbf{s}(\mathbf{a}^m, \mathbf{b}_m) = (s_1, s_2, \dots, s_T)$. Then it follows from (2) that $\mathbf{s}(\mathbf{a}^{m-1}, \mathbf{b}_{m-1}) = (s_2, s_3, \dots, s_T)$. Hence,

$$q_1 = m - 1 - \sum_{t=2}^T (t-1) s_t = m - 1 - \sum_{t=1}^T t s_t + \sum_{t=1}^T s_t = q - 1 + \sum_{t=1}^T s_t.$$

From (3), we have $q_2 - q_1 = \sum_{t=2}^T s_t - 1 \leq \sum_{t=1}^T s_t - 1 = q_1 - q$. This proves (4). ■

Applying Lemma 3.10 to the quantities in Definition 3.2, we readily deduce the following.

Lemma 3.11 *Use the notation in Definition 3.2 and 3.9. The following conditions hold.*

- (1) $k = T(\mathbf{b}_{n-p_0}, \mathbf{a}_{n-p_0})$, $\ell = T(\mathbf{a}_{n-q_0}, \mathbf{b}_{n-q_0})$.
(2) Suppose $\mathbf{s}(\mathbf{b}_{n-p_0}, \mathbf{a}_{n-p_0}) = (s_1, s_2, \dots, s_k)$ and $\mathbf{s}(\mathbf{a}_{n-q_0}, \mathbf{b}_{n-q_0}) = (s'_1, s'_2, \dots, s'_\ell)$.
Then

$$\begin{aligned} Q_{i+1} &= Q_i - 1 + \sum_{t=i+1}^k s_t && \text{for } 0 \leq i < k, \\ P_{j+1} &= P_j - 1 + \sum_{t=j+1}^{\ell} s'_t && \text{for } 0 \leq j < \ell. \end{aligned}$$

- (3) For $k \leq i < n - (p_0 + q_0 + \ell)$ and $\ell \leq j < n - (p_0 + q_0 + k)$, we have

$$Q_{i+1} = Q_i - 1 \quad \text{and} \quad P_{j+1} = P_j - 1.$$

Moreover, for $k \leq i \leq n - (p_0 + q_0 + \ell)$ and $\ell \leq j \leq n - (p_0 + q_0 + k)$, we have

$$p_i + Q_i = n = P_j + q_j. \tag{3.10}$$

- (4) For $0 < i < n - (p_0 + q_0 + \ell)$ and $\ell < j < n - (p_0 + q_0 + k)$, we have

$$Q_i - Q_{i-1} \geq Q_{i+1} - Q_i \quad \text{and} \quad P_j - P_{j-1} \geq P_{j+1} \geq P_{j+1} - P_j$$

Proof of Theorem 3.3 (1) From (p_0, q_0) to (p_0, Q_0) , we have a vertical straight line segment. Note that the slope of the line segment from (p_{i-1}, Q_{i-1}) to (p_i, Q_i) equals $Q_i - Q_{i-1}$, and the slope of the line segment from (p_i, Q_i) to (p_{i+1}, Q_{i+1}) is $Q_{i+1} - Q_i$. By Lemma 3.11 (4), we see that $Q_i - Q_{i-1} \geq Q_{i+1} - Q_i$. Thus, the polygonal curve joining the points $(p_0, Q_0), (p_1, Q_1), \dots, (p_k, Q_k)$ is convex. The line segment joining (p_k, Q_k) and (P_ℓ, q_ℓ) is a line segment with negative slope. Finally, the polygonal curve joining the points $(p_0, q_0), (P_0, q_0), \dots, (P_\ell, q_\ell)$ is concave by Lemma 3.11 (4). Thus \mathcal{P} is a convex subset contained in the set

$$\{(p, q) : p_0 \leq p \leq n - q_\ell, q_0 \leq q \leq n - p_k, \text{ and } p + q \leq n\}.$$

(2) Suppose $(p, q) \in \mathcal{P}$. Let $p = p_i$ and $q = q_j$ for some $0 \leq i \leq n - (p_0 + q_0 + \ell)$ and $0 \leq j \leq n - (p_0 + q_0 + k)$. Then $p_i \leq P_j$ and $q_j \leq Q_i$. Since (p_i, Q_i) and $(P_j, q_j) \in \text{In}(\mathbf{a}, \mathbf{b})$. By Corollary 2.7, $(p_i, q_j) \in \text{In}(\mathbf{a}, \mathbf{b})$.

Conversely, suppose $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$. By Theorem 3.6, we have $p \geq p_0$, $q \geq q_0$ and $p + q \leq n$. Let $p = p_i$ and $q = q_j$ for some $i, j \geq 0$. If $i > n - (p_0 + q_0 + \ell)$, then we have

$$q_j \leq n - p_i < q_0 + \ell \Rightarrow p_i \leq P_j \leq n - q_\ell \Rightarrow i \leq n - (p_0 + q_0 + \ell),$$

a contradiction. Therefore, $0 \leq i \leq n - (p_0 + q_0 + \ell)$. Similarly, we have $0 \leq j \leq n - (p_0 + q_0 + k)$. Since $(p_i, q_j) \in \text{In}(\mathbf{a}, \mathbf{b})$, we have $p_i \leq P_j$ and $q_j \leq Q_i$ by Lemma 3.8. If either $p_i = P_j$ or $q_j = Q_i$, then $(p, q) \in \mathcal{P}$. So we may assume that $p_i < P_j$ and $q_j < Q_i$. Consider the positive numbers

$$t_1 = j(P_j - p_i), \quad t_2 = i(Q_i - q_j) \quad \text{and} \quad t_3 = (P_j - p_i)(Q_i - q_j).$$

Then, by direct computation, we have

$$\frac{t_1(p_i, Q_i) + t_2(P_j, q_j) + t_3(p_0, q_0)}{t_1 + t_2 + t_3} = \frac{(t_1 p_i + t_2 P_j + t_3 p_0, t_1 Q_i + t_2 q_j + t_3 q_0)}{t_1 + t_2 + t_3} = (p_i, q_j).$$

Thus, (p, q) lies in \mathcal{P} . ■

4 Elements in $\text{In}(\mathbf{a}, \mathbf{b})$ attainable by diagonal matrices

In this section, we determine those elements in $\text{In}(\mathbf{a}, \mathbf{b})$ that are attainable by diagonal matrices. Clearly, if A and B are diagonal matrices with eigenvalues so that the eigenvalues of A and those of B are mutually distinct, then $A - B$ is invertible. If A and B have m common eigenvalues (counting multiplicities), then $A - B$ has at most m zero eigenvalues. It turns out that this is the only additional restriction on $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ to be attainable by diagonal matrices.

Theorem 4.1 *Suppose \mathbf{a} and \mathbf{b} have m common entries counting multiplicities. Then there are diagonal matrices $A \in \mathcal{H}_n(\mathbf{a})$ and $B \in \mathcal{H}_n(\mathbf{b})$ such that $A - B \in \mathcal{H}_n(p, q)$ if and only if $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ and $p + q \geq n - m$.*

To prove Theorem 4.1 we need the following.

Lemma 4.2 *Let $\mathbf{a} = (a_1, a_2, \dots, a_n)$ and $\mathbf{b} = (b_1, b_2, \dots, b_n) \in \mathbf{R}^n$ with $a_1 \geq a_2 \geq \dots \geq a_n$ and $b_1 \geq b_2 \geq \dots \geq b_n$. Given $1 \leq j_1 \leq i_1 \leq n$, let $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ be obtained from \mathbf{a} and \mathbf{b} by deleting a_{i_1} and b_{j_1} from \mathbf{a} and \mathbf{b} respectively. Suppose $\mathbf{a} \succ_p \mathbf{b}$ for some $0 \leq p \leq n$. We have*

- (1) $\hat{\mathbf{a}} \geq \hat{\mathbf{b}}$.
- (2) If $1 \leq p \leq n$, then $\hat{\mathbf{a}} \geq_{p-1} \hat{\mathbf{b}}$.
- (3) If $a_i = b_i$ for some $j_1 \leq i \leq i_1$, then $\hat{\mathbf{a}} \geq_p \hat{\mathbf{b}}$.

Proof. Since

$$\hat{a}_i = \begin{cases} a_i & \text{if } 1 \leq i < i_1, \\ a_{i+1} & \text{if } i_1 \leq i \leq n-1, \end{cases} \quad \text{and} \quad \hat{b}_j = \begin{cases} b_j & \text{if } 1 \leq j < j_1, \\ b_{j+1} & \text{if } j_1 \leq j \leq n-1, \end{cases}$$

we have

$$\begin{aligned} 1 \leq i < j_1 &\Rightarrow \hat{a}_i = a_i \geq b_i = \hat{b}_i \\ j_1 \leq i < i_1 &\Rightarrow \hat{a}_i = a_i \geq b_i \geq b_{i+1} = \hat{b}_i \\ i_1 \leq i < n &\Rightarrow \hat{a}_i = a_{i+1} \geq b_{i+1} = \hat{b}_i \end{aligned} \tag{4.1}$$

and (1) holds.

Note that every strict inequality $a_i > b_i$ for $1 \leq i < i_1$ (or $i_1 < i \leq n$) gives a strict inequality $\hat{a}_i > \hat{b}_i$ (or $\hat{a}_{i-1} > \hat{b}_{i-1}$). This proves (2) and the case when $i = i_1$ or j_1 in (3).

For (3), we may assume that $a_{i_1} > b_{i_1}$ and $i_1 > j_1$. Note that

$$\begin{aligned} (\hat{a}_1, \hat{a}_2, \dots, \hat{a}_{j_1-1}) &= (a_1, a_2, \dots, a_{j_1-1}) \\ (\hat{b}_1, \hat{b}_2, \dots, \hat{b}_{j_1-1}) &= (b_1, b_2, \dots, b_{j_1-1}) \\ (\hat{a}_{j_1}, \hat{a}_{j_1+1}, \dots, \hat{a}_{i_1-1}) &= (a_{j_1}, a_{j_1+1}, \dots, a_{i_1-1}) \\ (\hat{b}_{j_1}, \hat{b}_{j_1+1}, \dots, \hat{b}_{i_1-1}) &= (b_{j_1+1}, b_{j_1+2}, \dots, b_{i_1}) \\ (\hat{a}_{i_1}, \hat{a}_{i_1+1}, \dots, \hat{a}_{n-1}) &= (a_{i_1+1}, a_{i_1+2}, \dots, a_n) \\ (\hat{b}_{i_1}, \hat{b}_{i_1+1}, \dots, \hat{b}_{n-1}) &= (b_{i_1+1}, b_{i_1+2}, \dots, a_n). \end{aligned}$$

Apply Lemma 3.10 (3) to $(a_{j_1}, a_{j_1+1}, \dots, a_{i_1})$ and $(b_{j_1}, b_{j_1+2}, \dots, b_{i_1})$; by the fact that at least one s_k is positive, we can conclude that the number of strict inequalities in $(\hat{a}_{j_1}, \hat{a}_{j_1+1}, \dots, \hat{a}_{i_1-1}) - (\hat{b}_{j_1}, \hat{b}_{j_1+1}, \dots, \hat{b}_{i_1-1})$ is no less than that of $(a_{j_1}, a_{j_1+1}, \dots, a_{i_1}) - (b_{j_1}, b_{j_1+2}, \dots, b_{i_1})$. Therefore, the number of entries in $\hat{\mathbf{a}} - \hat{\mathbf{b}}$ is no less than that of $\mathbf{a} - \mathbf{b}$.

■

Proof of Theorem 4.1. Suppose A and B are diagonal matrices with eigenvalues a_1, \dots, a_n and b_1, \dots, b_n such that $A - B \in \mathcal{H}_n(p, q)$. So, $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$. Also, the number of zero diagonal entries is at most m . Therefore, $m \geq n - p - q$. Hence, $p + q \geq n - m$.

We prove the converse by induction on m . Let $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ and $p + q \geq n - m$. If $p + q = n$ then the result follows from Theorem 2.1. So the result holds for $m = 0$ and we may assume that $n > p + q$.

Let $m > 0$. Assume the result holds whenever \mathbf{a} and \mathbf{b} have $m - 1$ entries in common. Suppose \mathbf{a} and \mathbf{b} have m common entries and $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$, with $p + q \geq n - m$. By Theorem 2.1, we have $\mathbf{a}^{n-q} \geq_p \mathbf{b}_{n-q}$ and $\mathbf{b}^{n-p} \geq_q \mathbf{a}_{n-p}$. We may assume that $n > p + q \geq n - m$. We are going to show that we can delete a common entries from \mathbf{a} and \mathbf{b} to obtain vectors $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}} \in \mathbf{R}^{n-1}$ so that $\hat{\mathbf{a}}^{n-1-q} \geq_p \hat{\mathbf{b}}_{n-1-q}$ and $\hat{\mathbf{b}}^{n-1-p} \geq_q \hat{\mathbf{a}}_{n-1-p}$. Since $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ have only $m - 1$ entries in common and $p + q \geq (n - 1) - (m - 1)$, the result will follow.

Consider the following cases:

Case 1: $\mathbf{a}^{n-q} \geq_{p+1} \mathbf{b}_{n-q}$ and $\mathbf{b}^{n-p} \geq_{q+1} \mathbf{a}_{n-p}$.

Since $m > 0$, we can choose $i_1 = \min\{i : a_i = b_j \text{ for some } j\}$ and $j_1 = \min\{j : b_j = a_{i_1}\}$. Let $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ be obtained from \mathbf{a} and \mathbf{b} by deleting a_{i_1} and b_{j_1} respectively.

If $i_1 > n - q$, then $\hat{\mathbf{a}}^{n-1-q} = \mathbf{a}^{n-1-q}$. Therefore, $\hat{\mathbf{a}}^{n-1-q} \geq_p \hat{\mathbf{b}}_{n-1-q}$.

If $i_1 \leq n - q$, then $b_{j_1-1} > b_{j_1} = a_{i_1} \geq b_{q+i_1}$ and we have $q + i_1 \geq j_1$. By Lemma 4.2 (2), $\hat{\mathbf{a}}^{n-1-q} \geq_p \hat{\mathbf{b}}_{n-1-q}$.

Similarly, we have $\hat{\mathbf{b}}^{n-1-p} \geq_q \hat{\mathbf{a}}_{n-1-p}$.

Case 2: $\mathbf{a}^{n-q} >_p \mathbf{b}_{n-q}$.

Since $n - q > p$, let $i_1 = \min\{t : 1 \leq t \leq n - q \text{ and } a_t = b_{q+t}\} \leq p + 1$. Let $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ be obtained from \mathbf{a} and \mathbf{b} by deleting a_{i_1} and b_{q+i_1} respectively. Then $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ have $m - 1$ entries in common. By Lemma 4.2 (3), $\hat{\mathbf{a}}^{n-1-q} \geq_p \hat{\mathbf{b}}_{n-1-q}$. Consider the following cases:

Subcase 2a: If $\mathbf{b}^{n-p} \geq_{q+1} \mathbf{a}_{n-p}$, then it follow from Lemma 4.2 (2) that $\hat{\mathbf{b}}^{n-1-p} \geq_q \hat{\mathbf{a}}_{n-1-p}$.

Subcase 2b: If $\mathbf{b}^{n-p} >_q \mathbf{a}_{n-p}$, then

$$\min\{s : 1 \leq s \leq n - p \text{ and } b_s = a_{p+s}\} \leq q + 1 \leq q + i_1.$$

It follow from Lemma 4.2 (3) that $\hat{\mathbf{b}}^{n-1-p} \geq_q \hat{\mathbf{a}}_{n-1-p}$. ■

5 Ranks and multiple eigenvalues

By Theorem 3.3, we can determine the set $R(\mathbf{a}, \mathbf{b})$ of all possible ranks a matrix of the form $A - B$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$. Evidently, we have

$$R(\mathbf{a}, \mathbf{b}) = \{p + q : (p, q) \in \text{In}(\mathbf{a}, \mathbf{b})\}.$$

Nevertheless, it is interesting that the result can be put in the following simple form.

Theorem 5.1 *Let \mathbf{a}, \mathbf{b} be real vectors, and define p_0 and q_0 as in (3.1) and (3.2). Let m be the largest multiplicity of an entry in $(a_1, \dots, a_n, b_1, \dots, b_n)$ and $r = \min\{2n - m, n\}$. Suppose $R(\mathbf{a}, \mathbf{b})$ is the set of rank values of matrices of the form $A - B$, where $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$. Then one of the following holds.*

(1) *There exist real numbers $\mu > \nu$ and $0 \leq u, v \leq n$ such that*

$$\mu = a_1 = \dots = a_u = b_1 = \dots = b_v, \quad \nu = a_{u+1} = \dots = a_n = b_{v+1} = \dots = b_n,$$

and

$$R(\mathbf{a}, \mathbf{b}) = \{u + v - 2j : \max\{0, u + v - n\} \leq j \leq \min\{u, v\}\}.$$

(2) Condition (1) does not hold, $\mathbf{a} = \mathbf{b}$, and

$$R(\mathbf{a}, \mathbf{b}) = \{0\} \cup \{2, \dots, r\}.$$

(3) Conditions (1) and (2) do not hold, and

$$R(\mathbf{a}, \mathbf{b}) = \{p_0 + q_0, \dots, r\}.$$

Moreover, if $t \in R(\mathbf{a}, \mathbf{b})$ then there are block diagonal matrices $A = A_1 \oplus \dots \oplus A_t \in \mathcal{H}_n(\mathbf{a})$ and $B = B_1 \oplus \dots \oplus B_t$ in $\mathcal{H}_n(\mathbf{b})$ with the same block sizes such that $A_j - B_j$ has rank one for $j = 1, \dots, t$.

Note that in the theorem, we include the case when $(a_1, \dots, a_n, b_1, \dots, b_n)$ has an entry with multiplicity larger than n .

Proof. (1) Suppose \mathbf{a}, \mathbf{b} satisfy the condition in (1). The result follows from Corollary 3.5.

(2) Suppose condition (1) does not hold and $\mathbf{a} = \mathbf{b}$. If $A = B = \text{diag}(a_1, \dots, a_n)$, then $A - B \in \mathcal{H}_n(0, 0)$. Since A and B have the same trace, we see that $A - B$ cannot have rank 1.

Without loss of generality, we may assume that $r = n$. We prove the following claim by induction on n :

There are matrices $A, B \in \mathcal{H}_n(\mathbf{a})$ such that $A - B \in \mathcal{H}_n(p, q)$ whenever $2 \leq p + q \leq n$ with $p = q$ or $p = q + 1$.

The claim is clear if $n = 3, 4$. Suppose $n \geq 5$ and $2 \leq p + q \leq n$ with $p = q$ or $p = q + 1$. Since \mathbf{a} has at least three distinct entries, each entry has multiplicity at most $n/2$. Suppose $a_r > a_s$, where a_r, a_s have the two largest multiplicities in the vector \mathbf{a} .

For $2 \leq p + q \leq 3$, choose $a_w \notin \{a_u, a_v\}$ and let $A_1 = \text{diag}(a_u, a_v, a_w)$. Then there exists a diagonal matrix B_1 with the same eigenvalues as A_1 and $A_1 - B_1 \in \mathcal{H}_3(p, q)$. Remove a_u, a_v, a_w from \mathbf{a} to get \mathbf{a}' . Then $A_1 \oplus \text{diag}(\mathbf{a}') - B_1 \oplus \text{diag}(\mathbf{a}') \in \mathcal{H}_n(p, q)$.

For $4 \leq p + q \leq n$, we have $p, q \geq 1$. Therefore, $2 \leq (p - 1) + (q - 1) \leq n - 2$ and $p - 1 = q - 1$ or $p - 1 = (q - 1) + 1$. Let $A_1 = \text{diag}(a_u, a_v)$ and $B_1 = \text{diag}(a_v, a_u)$, we have $A_1 - B_1 \in \text{In}(1, 1)$. Remove a_r, a_s from \mathbf{a} to get \mathbf{a}' . Since $n \geq 5$, there are at least three distinct entries in \mathbf{a}' and each has multiplicity at most $(n - 2)/2$. By induction assumption, there are A_2, B_2 both with vector of eigenvalues \mathbf{a}' such that $A_2 - B_2 \in \mathcal{H}_{n-2}(p - 1, q - 1)$. Thus, $A_1 \oplus A_2 - B_1 \oplus B_2 \in \mathcal{H}_n(p, q)$.

(3) Suppose conditions (1) and (2) do not hold. Using the notation in Theorem 3.3, we see that $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ for

$$(p, q) \in \{(p_j, q_0) : 0 \leq j \leq k\} \cup \{(p_k, q_j) : 1 \leq j \leq Q_k\}.$$

Thus, we have the desired rank values.

By Theorem 2.1, we can construct matrices A and B with the asserted block structure. ■

It is clear that $X, Y \in \mathcal{H}_n$ have the same eigenvalues if and only if $X - \mu I$ and $Y - \mu I$ have the same inertia (or rank) for all eigenvalues μ of Y . Thus, we can describe the eigenvalues of $A - B$ in terms of the inertia of $A - B - \mu I$ for different real numbers μ . In particular, we have the following necessary condition for $c_1 \geq \dots \geq c_n$ to be the eigenvalues of $A - B$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$.

Proposition 5.2 *Let $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n)$, $\mathbf{c} = (c_1, \dots, c_n)$ be real vectors with entries arranged in descending order. Suppose \mathbf{c} has distinct entries $c_1 > \dots > c_t$ with multiplicities m_1, \dots, m_t , respectively, and suppose there exists $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ such that $A - B \in \mathcal{H}_n(\mathbf{c})$. Set $u_0 = 0$, $u_j = m_1 + \dots + m_{j-1}$ for $j \in \{1, \dots, t\}$, $v_j = m_{j+1} + \dots + m_t$ for $j \in \{1, \dots, t-1\}$ and $v_t = 0$. Then for $j \in \{1, \dots, t\}$,*

- (i) $(a_1 - c_j, \dots, a_{n-v_j} - c_j) - (b_{v_j+1}, \dots, b_n)$ is nonnegative with at least u_j positive entries.
- (ii) $(b_1, \dots, b_{n-u_j}) - (a_{u_j+1} - c_j, \dots, a_n - c_j)$ is nonnegative with at least v_j positive entries.

Remark 5.3 Let $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ with entries arranged in descending order. Then there exist $A, B \in \mathcal{H}_n$ with vector of eigenvalues \mathbf{a} and \mathbf{b} such that $A - B$ has an eigenvalue μ with multiplicity t if and only if there is a matrix of the form $\tilde{A} - B$ has rank $n - t$, where $\tilde{A} + \mu I \in \mathcal{H}_n(\mathbf{a})$ and $B \in \mathcal{H}_n(\mathbf{b})$. Hence, we can use Theorem 5.1 to determine whether there is $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ such that $A - B$ has an eigenvalue μ with multiplicity t . In Corollaries 5.6 and 5.7, we will apply Theorem 2.1 to give a more precise location of the multiple eigenvalue μ . As a byproduct, we determine the function $f(\mu)$ defined as the minimum rank of a matrix of the form $A - B - \mu I$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ for given real vectors \mathbf{a} and \mathbf{b} .

The following notation will be used for the rest of this section.

Notation 5.4 *Let $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n)$ be real vectors with entries arranged in descending order. For $0 \leq t \leq n - 1$, let*

$$\alpha_t = \max\{a_{j+t} - b_j : 1 \leq j \leq n - t\} \quad \text{and} \quad \beta_t = \min\{a_j - b_{j+t} : 1 \leq j \leq n - t\}.$$

For $\mu \in \mathbf{R}$, let $p_0(\mu)$ and $q_0(\mu)$ be defined as in (3.1) – (3.2), with a_j replaced by $a_j - \mu$.

Note that $p_0(\mu) + q_0(\mu)$ will be the minimum rank of a matrix of the form $A - B - \mu I$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$.

Proposition 5.5 *Let \mathbf{a} and \mathbf{b} be real vectors with entries arranged in descending order. We have*

$$\alpha_{n-1} \leq \alpha_{n-2} \leq \dots \leq \alpha_0 \quad \text{and} \quad \beta_0 \leq \beta_1 \leq \dots \leq \beta_{n-1}.$$

Moreover, the following conditions hold for the function $p_0(\mu)$, $q_0(\mu)$ and $p_0(\mu) + q_0(\mu)$.

- (a) $p_0(\mu)$ is a decreasing step function in $\mu \in \mathbf{R}$ such that $p_0(\mu) = n$ for $\mu < \alpha_{n-1}$, $p_0(\mu) = 0$ for $\mu \geq \alpha_0$, and $p_0(\mu) = t$ if μ in the interval $[\alpha_t, \alpha_{t-1})$ for $1 \leq t \leq n-1$;
- (b) $q_0(\mu)$ is an increasing step function in $\mu \in \mathbf{R}$ such that $q_0(\mu) = 0$ for $\mu \leq \beta_0$, $q_0(\mu) = n$ for $\mu > \beta_{n-1}$, and $q_0(\mu) = t$ if μ in the interval $(\beta_{t-1}, \beta_t]$ for $1 \leq t \leq n-1$.
- (c) If $\alpha_s = \beta_t$ for some $0 \leq s, t \leq n-1$, then there exists $\delta > 0$ such that $p_0(\mu) + q_0(\mu) > p_0(\alpha_s) + q_0(\alpha_s)$ for all $0 < |\mu - \alpha_s| < \delta$.
- (d) If $\mu \neq \alpha_t, \beta_t$ for all $0 \leq t \leq n-1$, then $p_0(\cdot) + q_0(\cdot)$ is locally constant at μ .

Proof. For $1 \leq t \leq n-1$ and $1 \leq j \leq n-t$ we have $a_{j+t} - b_j \leq a_{j+(t-1)} - b_j$. Therefore, $\alpha_t \leq \alpha_{t-1}$. Similarly, $\beta_t \geq \beta_{t-1}$.

By (3.1) and (3.2), we have

$$p_0(\mu) = \begin{cases} n & \text{if } \mu < a_n - b_1, \\ \min\{t : \mu \geq \alpha_t\} & \text{otherwise,} \end{cases}$$

$$q_0(\mu) = \begin{cases} n & \text{if } a_1 - b_n < \mu, \\ \min\{t : \mu \leq \beta_t\} & \text{otherwise,} \end{cases}$$

which implies (a) and (b).

For (c), suppose $\alpha_s = \beta_t$ for some $0 \leq s, t \leq n-1$. By taking $\alpha_n = \alpha_{n-1} - 1$, $\alpha_{-1} = \alpha_0 + 1$, $\beta_{-1} = \beta_0 - 1$ and $\beta_n = \beta_{n-1} + 1$, we may assume that $\alpha_{s+1} < \alpha_s = c_{s-1} = \dots = \alpha_{s'} < \alpha_{s'-1}$ and $\beta_{t-1} < \beta_t = \beta_{t+1} = \dots = \beta_{t'} < \beta_{t'+1}$. Let $\delta = \min\{\alpha_s - \alpha_{s+1}, \alpha_{s'-1} - \alpha_{s'}, \beta_t - \beta_{t-1}, \beta_{t'+1} - \beta_{t'}\} > 0$. We have $p_0(\mu) + q_0(\mu) = s + t + 1 > s' + t = p_0(\alpha_{s'}) + q_0(\beta_t) = p_0(\alpha_s) + q_0(\alpha_s)$ if $0 < \alpha_s - \mu < \delta$ and $p_0(\mu) + q_0(\mu) = s' + t' + 1 > s' + t$ if $0 < \mu - \alpha_s < \delta$.

$$p_0(\mu) + q_0(\mu) = \begin{cases} s + t + 1 & \text{if } 0 < \alpha_s - \mu < \delta \\ s' + t' + 1 & \text{if } 0 < \mu - \alpha_s < \delta \end{cases}$$

$$> s' + t = p_0(\alpha_{s'}) + q_0(\beta_t) = p_0(\alpha_s) + q_0(\alpha_s)$$

(d) follows from (a) and (b). ■

Note that the function $g(\mu)$ defined as the maximum rank of a matrix of the form $A - B - \mu I$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ is easy to determine, namely, it is equal to $g(\mu) = \min\{n, 2n - m(\mu)\}$ with $m(\mu)$ equal to the maximum multiplicity of an entry in the vector $(a_1 - \mu, \dots, a_n - \mu, b_1, \dots, b_n)$.

Similarly, one can consider $P_\ell(\mu)$ and $Q_k(\mu)$ defined as the maximum number of positive and negative eigenvalues of a matrix of the form $A - B - \mu I$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}(\mathbf{b})$. We omit their discussion.

The following corollary concerns the possible multiplicities for $\mu \in \mathbf{R}$ to be an eigenvalue of $A - B$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$.

Corollary 5.6 *Let \mathbf{a} and \mathbf{b} be real vectors with entries arranged in descending order. Suppose $a_n - b_1 \leq \mu \leq a_1 - b_n$. Then there exist $s, t \in \{0, \dots, n-1\}$ such that $\mu \in [\alpha_s, \alpha_{s-1}) \cap (\beta_{t-1}, \beta_t]$, where we take $\alpha_{-1} > \beta_{n-1}$ and $\beta_{-1} < \alpha_{n-1}$.*

- (1) *Suppose $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ and μ is an eigenvalue of $A - B$. Then the multiplicity of μ is at most $n - s - t$. Furthermore, $A - B$ has at least s eigenvalue greater than μ and at least t eigenvalues less than μ .*
- (2) *There exists $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ such that $A - B$ has an eigenvalue μ with multiplicity $n - s - t$, s eigenvalues greater than μ and t eigenvalues less than μ .*

To facilitate the comparison of our results and those in the literature, we present the next corollary in terms of $A + B$ with $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$. We use the following notation. Let $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n)$ and $\mathbf{c} = (c_1, \dots, c_n)$ with entries arranged in descending order. For each $1 \leq k \leq n$, let $L_k = \max\{a_i + b_j : i + j = n + k\}$ and $R_k = \min\{a_i + b_j : i + j = k + 1\}$. Suppose $A, B \in \mathcal{H}_n$ and $C = A + B$ have eigenvalues \mathbf{a} , \mathbf{b} and \mathbf{c} . Then it follows from Weyl's inequalities [13] that $L_k \leq c_k \leq R_k$. Conversely, for every $1 \leq k \leq n$ and $c \in [L_k, R_k]$, there exist $A, B \in \mathcal{H}_n$ and $C = A + B$ with eigenvalues \mathbf{a} , \mathbf{b} and \mathbf{c} such that $c_k = c$. However, for two distinct $1 \leq k < k' \leq n$ and $c \in [L_k, R_k]$, $c \in [L_{k'}, R_{k'}]$, there may not exist $A, B \in \mathcal{H}_n$ and $C = A + B$ with eigenvalues \mathbf{a} , \mathbf{b} and \mathbf{c} such that $c_k = c$ and $c_{k'} = c'$; see the example in [7, p.215]. Nevertheless, by replacing b_j with $-b_{n+1-j}$ and putting $s = k - 1$ and $t = n - k'$, the second part of Corollary 5.6 can be rephrased in the following form.

Corollary 5.7 *Let $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ with entries arranged in descending order and $\mu \in [L_k, L_{k-1}) \cap (R_{k'+1}, R_{k'}]$. Then there exists $(A, B) \in \mathcal{H}_n(\mathbf{a}, \mathbf{b})$ such that $C = A + B$ has a vector of eigenvalues \mathbf{c} with $c_{k-1} < \mu = c_k = c_{k+1} = \dots = c_{k'} < c_{k'+1}$.*

We remark that Corollary 5.7 can also be deduced from the results in [1].

6 Additional results and remarks

Proposition 6.1 *Let \mathbf{a}, \mathbf{b} be given. There are $1 \times n$ vectors \mathbf{a}' and \mathbf{b}' with integral entries arranged in descending order such that $\text{In}(\mathbf{a}, \mathbf{b}) = \text{In}(\mathbf{a}', \mathbf{b}')$. Moreover, for each $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ there is $A \in \mathcal{H}_n(\mathbf{a}')$ and $B \in \mathcal{H}_n(\mathbf{b}')$ such that $A - B \in \text{In}(\mathbf{a}', \mathbf{b}')$ has integer eigenvalues.*

Proof. We can construct \mathbf{a}' and \mathbf{b}' as follows. Use the entries of \mathbf{a} and \mathbf{b} to form a vector $\gamma = (\gamma_1, \dots, \gamma_{2n})$ with entries in descending order. We always put the entries of \mathbf{a} first if an entry appears in both vectors. Suppose γ has m distinct entries $\mu_1 > \dots > \mu_m$. Then replace the entries μ_i in \mathbf{a} and \mathbf{b} by the integer i for each $i \in \{1, \dots, m\}$ to get the vectors \mathbf{a}' and \mathbf{b}' . By Theorem 2.1 and the construction of \mathbf{a}' and \mathbf{b}' , we see that $(p, q) \in \text{In}(\mathbf{a}, \mathbf{b})$ if and only if $(p, q) \in \text{In}(\mathbf{a}', \mathbf{b}')$. Moreover, by Theorem 2.1, for each $(p, q) \in \text{In}(\mathbf{a}', \mathbf{b}')$ we can construct $A = A_1 \oplus \dots \oplus A_{p+q} \in \mathcal{H}_n(\mathbf{a}')$ and $B = B_1 \oplus \dots \oplus B_{p+q} \in \mathcal{H}_n(\mathbf{b}')$ such that $A_i - B_i$ is a rank one positive semi-definite for $i = 1, \dots, p$, and $A_i - B_i$ is a rank one negative semi-definite for $i = p + 1, \dots, p + q$. Since A_i and B_i has integral eigenvalues, the only nonzero eigenvalue of $A_i - B_i$ equals $\text{tr}(A_i - B_i)$ is again an integer. So, the last assertion holds. \blacksquare

Suppose $\mathbf{a}, \mathbf{b}, \mathbf{c}$ have nonnegative integral entries. It is known that there exist $(A, B) \in \mathcal{H}_n(\mathbf{a}) \times \mathcal{H}_n(\mathbf{b})$ such that $A - B \in \mathcal{H}_n(\mathbf{c})$ if and only if one can obtain the Young diagram associated with (a_1, \dots, a_n) from the Young diagrams associated with (b_1, \dots, b_n) and (c_1, \dots, c_n) according to the Little-Richardson rules; see [7]. Thus, we have the following result.

Proposition 6.2 *Let $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ have positive integral entries arranged in descending order. Then there is a vector $\mathbf{c} = (c_1, \dots, c_n)$ with positive integral entries arranged in descending order and $c_{p+1} = \dots = c_{n-q+1} = \mu$ for a given integer μ such that one can obtain the Young diagram associated with \mathbf{a} from the Young diagrams associated with \mathbf{b} and \mathbf{c} according to the Little-Richardson rules if and only if*

$$(a_1 - \mu, \dots, a_{n-q} - \mu) \geq_p (b_{q+1}, \dots, b_n)$$

and

$$(b_1, \dots, b_{n-p}) \geq_q (a_{p+1} - \mu, \dots, a_n - \mu).$$

In connection to our discussion, it would be interesting to solve the following.

Problem 6.3 *Deduce and extend Proposition 6.2 using the theory of algebraic combinatorics. In particular, for given real vectors \mathbf{a} and \mathbf{b} with integral entries, determine the conditions for the existence of an integral vectors \mathbf{c} with certain prescribed entries such that the Young diagram corresponding to \mathbf{a} can be obtained from those of \mathbf{b} and \mathbf{c} according to the Littlewood-Richardson rules.*

Problem 6.4 *Extend our results to the sum of k Hermitian matrices for $k > 2$. In other words, determine all possible inertia values and ranks of matrices in $\mathcal{H}_n(\mathbf{a}_1) + \dots + \mathcal{H}_n(\mathbf{a}_k)$ for given $1 \times n$ real vectors $\mathbf{a}_1, \dots, \mathbf{a}_k$ with entries arranged in descending order.*

Note that the problem of finding the relation between the eigenvalues of A_1, \dots, A_k and that of $A_0 = A_1 + \dots + A_k$ can be reformulated as the problem of finding the necessary and sufficient conditions for the existence of Hermitian matrices A_0, A_1, \dots, A_k with prescribed eigenvalues such that $A_0 - \sum_{j=1}^k A_j$ has rank 0. Thus, it can be viewed as a special case of Problem 6.4. To determine whether there are $A_1, \dots, A_k \in \mathcal{H}_n$ with prescribed eigenvalues such that $A_1 + \dots + A_k$ has rank one, one may reduce the problem to the study of the existence of $A_1, \dots, A_k \in \mathcal{H}_n$ with prescribed eigenvalues such that $A_1 + \dots + A_k$ has eigenvalue $\mu, 0, \dots, 0$ with $\mu = \text{tr}(A_1 + \dots + A_k)$. Then the results in [7] can be used to solve the problem. In general, it seems difficult to determine if there exist A_1, \dots, A_k with prescribed eigenvalues such that $A_1 + \dots + A_k$ has rank r with $r \in \{2, \dots, n\}$.

Note added in proof.

We thank Professor Wing Suet Li for some helpful discussion about the connection of the interesting preprint [1] and our work. In [1, Proposition 5.1], the authors determined the conditions on $1 \times n$ vectors $\mathbf{a}_0, \mathbf{a}_1, \dots, \mathbf{a}_k$, with some of their entries specified so that one can fill in the missing entries to get vectors $\tilde{\mathbf{a}}_0, \dots, \tilde{\mathbf{a}}_k$ with entries arranged in descending order and Hermitian matrices $A_j \in \mathcal{H}_n(\tilde{\mathbf{a}}_j)$ for $j = 0, 1, \dots, k$ satisfying $A_0 = A_1 + \dots + A_k$. Evidently, there exists $A_0 \in \mathcal{H}(\mathbf{a}_1) + \dots + \mathcal{H}(\mathbf{a}_k)$ with inertia $(p, q, n - p - q)$ for given $1 \times n$ real vectors $\mathbf{a}_1, \dots, \mathbf{a}_k$ if and only if there exist $\varepsilon > 0$ and $A_0 \in \mathcal{H}(\mathbf{a}_1) + \dots + \mathcal{H}(\mathbf{a}_k)$ with eigenvalues $\mu_1 \geq \dots \geq \mu_n$ such that $(\mu_p, \dots, \mu_{n-q+1}) = (\varepsilon, 0, \dots, 0, -\varepsilon)$. Using the result in [1], one can determine whether the desired positive number ε exists by checking whether a polytope defined a large number of inequalities in terms of entries of $\mathbf{a}_1, \dots, \mathbf{a}_k$ has non-empty interior; see also Buch [2]. For $k = 2$, our Theorem 2.1 shows that the very involved conditions can be reduced to (1) and (2).

References

- [1] H. Bercovici, W.S. Li and D. Timotin, The Horn conjecture for sums of compact selfadjoint operators, preprint. <http://arxiv.org/abs/0709.1088>
- [2] A.S. Buch, Eigenvalues of Hermitian matrices with positive sum of bounded rank, *Linear Algebra Appl.* 418 (2006), 480-488.
- [3] M.D. Choi and P.Y. Wu, Convex combinations of projections, *Linear Algebra Appl.* 136 (1990), 25-42.
- [4] M.D. Choi and P.Y. Wu, Finite-rank perturbations of positive operators and isometries, *Studia Math.* 173 (2006), no. 1, 73-79.
- [5] J. Day, W. So and R.C. Thompson, The spectrum of a Hermitian matrix sum, *Linear Algebra Appl.* 280 (1998), no. 2-3, 289-332.

- [6] K. Fan and G. Pall, Imbedding conditions for Hermitian and normal matrices, *Canad. J. Math.* 9 (1957), 298-304.
- [7] W. Fulton, Eigenvalues, invariant factors, highest weights, and Schubert calculus, *Bull. Amer. Math. Soc.* 37 (2000), 209–249.
- [8] A. Horn, Eigenvalues of sums of Hermitian matrices, *Pacific J. Math* 12 (1962), 225–241.
- [9] A.A. Klyachko, Stable bundles, representation theory and Hermitian operators, *Selecta Math. (N.S.)* 4 (1998), 419-445.
- [10] A. Knutson and T. Tao, The honeycomb model of $GL_n(c)$ tensor products. I. Proof of the saturation conjecture, *J. Amer. Math. Soc.* 12 (1999), no. 4, 1055–1090.
- [11] R.C. Thompson and L.J. Freede, On the eigenvalues of sums of Hermitian matrices, *Linear Algebra and Appl.* 4 1971 369–376.
- [12] R.C. Thompson and L.J. Freede, On the eigenvalues of sums of Hermitian matrices, II. *Aequationes Math* 5 (1970), 103–115
- [13] H. Weyl, Das asymptotische Verteilungsgesetz der Eigenwerte lineare partieller Differentialgleichungen, *Math. Ann.* 71 (1912), 441–479

Department of Mathematics, College of William and Mary,
Williamsburg, VA 23185. (ckli@math.wm.edu)

Department of Mathematics, Iowa State University,
Ames, IA 50011. (ytpoon@iastate.edu)